

Sunrise C28

Sunrise C18

RP C18 column with feature of a silanol group

Sunrise C18-SAC

Silanol Activity Controlled C18 Column





Sunrise Octacosyl (C28)
Sunrise Octadecyl (C18)

Sunrise Octadecyl-SAC (C18-SAC) has an interaction of silanol groups

Sunrise C28

Sunrise C18



Name	Stationary phase	Carbon content	Ligand density	Particle size	
C18 Octadecyl		15%	2.1 $\mu\text{mol}/\text{m}^2$	3 μm , 5 μm	Silica support Surface area : 340 m^2/g Pore volume : 1.0 mL/g Pore diameter : 12 nm End-capping Trimethylsilyl group (TMS)
C28 Octacosyl		18%	1.7 $\mu\text{mol}/\text{m}^2$	3 μm , 5 μm	

pH range of C18 and dC28: pH2~pH8



◆ End-capping type Sunrise seriesの特徴

C28:

■ A long alkyl chain improves both separation of fat-soluble compounds to compare with C18 phase and an excellent reproducibility in retention under high aqueous conditions.

■ Furthermore, a suitable ligand density of C28 allows to be obtained a shape peak shape even if more than 50% aqueous mobile phase is used.

■ Different selectivity

C18:

■ Conventional C18 phase with full end-capping

PhE:


(Sunrise PhE has stopped production. Sunniest PhE is recommended as a replacement)

■ Interaction based with p-electron such as p-p interaction


■ p-electron also interacts with a polar site of a compound, so that phenyl phase improves separation of polar compounds. Ethylene chain between silica surface and phenyl group allows a movable sphere of a phenyl group to be wide. A chain with more than three carbons shows more hydrophobic interaction, so that p-electron interaction decreases relatively.

■ Phenethyl (PhE) group is a suitable phenyl phase.

■ Separation of Basic compounds



■ Separation of Vitamin E Isomer can be separated by C28




Sunrise C28


Sunrise C18



Comparison of stationary phases



Comparison of separation of α , β - carotene




The mobile phase including chloroform makes alkyl chains brush up because chloroform can enter among alkyl chains. Consequently retention times of C28 became 2 times longer than C18.


Sunrise C28



■ Separation of xylene isomers



■ Separation of ter-phenyl isomers



Sunrise C18-SAC


Silanol Activity Controlled C18 HPLC Column



◆ New generation reversed-phase utilized silanol groups


■ Silanol group and peak tailing

It is generally said that residual silanol groups on a stationary phase such as C18 (ODS) causes absorption or peak tailing for a sample. Especially silanol groups near a hydrophobic site don't solvate with water completely, so that they show high absorption for basic compounds. Its peak shows terribly tailing. Several end-capping techniques have been developed to solve these problems for many years.



■ Silanol activity control technology

ChromaNik developed the technique that decreased only silanol groups with high absorption activity to a basic compound and remained effective silanol groups on the stationary phase. Silanol activity control and no end-capping led the existence of silanol groups with high hydration which created a new and unique reversed-phase separation mode including hydrogen bond and ion-exchange interaction. Furthermore, silanol activity controlling, then end-capping technique improved a peak shape of a basic compound exceedingly.



◆ Feature of Sunrise series


Sunrise C18

- The "1st Choice" column as a fully end-capped C18 column
- Full end-capping after silanol activity control
- Reducing adsorption of a basic compound extremely
- A good peak shape for a metal chelating compound
- Widely available for general reversed-phase separation

Sunrise C18-SAC

- The "2nd Choice" column which takes advantage of effective silanol groups interaction
- Reducing silanol groups with high adsorption activity
- The new separation mechanism including hydrogen bond and ion-exchange interaction
- Effective for separation of a basic compound and a polar compound
- Different selectivity and improvement of separation without changing a mobile phase

■ The elution order of pyridine



Sunrise C18-SAC

Silanol Activity Controlled C18 HPLC Column



◆ Sunrise series create an unique separation


* Effectiveness of silanol activity control: Comparison between Sunrise C18 and C18-SAC

Sunrise C18 is the so-called fully end-capped C18 column. It shows the same separation behavior as a conventional C18 column.


On the other hand, Sunrise C18-SAC shows hydrogen bond and ion-exchange interactions based on a residual silanol on the silica support in addition to reversed-phase separation. For example Sunrise C18 column separates a basic compound similarly as a conventional C18, while

Sunrise C18-SAC makes retention of a basic compound be large because an ion-exchange interaction works although a non-ionic compound shows the almost same retention on both Sunrise C18 and C18-SAC. Furthermore, Sunrise C18-SAC shows large retention for a polar compound such as caffeine.


■ Comparison of selectivity for basic compounds



■ Comparison of peak shape and retention



■ Comparison of caffeine



Sunrise C18-SAC

Silanol Activity Controlled C18 HPLC Column




◆ Multiple mode separation is achieved on Sunrise series

* Silanol groups controlled its activity functions as ion-exchange groups


Sunrise C18-SAC is bonded with octadecylsilane on a pure silica gel and controlled its silanol activity without end-capping. Its carbon content is 14%.

Separation on Sunrise C18-SAC is done including hydrogen bond and ion-exchange interaction based on silanol groups except for hydrophobic interaction. Control of pH and salt concentration of a mobile phase can regulate retention.


■ Separation of basic compounds with ammonium acetate: Effect of salt concentration (Sunrise C18-SAC)



■ Relationship between buffer concentration and retention (Sunrise C18-SAC)



■ Chromatograms under different pH conditions (Sunrise C18-SAC)



Sunrise C28, C18, C18-SAC



* Sunrise series Analytical and Preparative Columns

Inner diameter [mm]	length [mm]	Sunrise C28, 3 μ m	Sunrise C28, 5 μ m
		Cat. No.	Cat. No.
2.0	50	ST2241	ST3241
	75	ST2251	—
	100	ST2261	ST3261
	150	ST2271	ST3271
	250	ST2281	ST3281
4.6	10	ST2411	ST3411
	50	ST2441	ST3441
	75	ST2451	—
	100	ST2461	ST3461
	150	ST2471	ST3471
	250	—	ST3481
10.0	250	—	ST3781
20.0	250	—	ST3881

Inner diameter [mm]	length [mm]	Sunrise C18, 3 μ m	Sunrise C18, 5 μ m	Sunrise C18-SAC, 3 μ m	Sunrise C18-SAC, 5 μ m
		Cat. No.	Cat. No.	Cat. No.	Cat. No.
2.0	50	SB2241	SB3241	SA2241	SA3241
	75	SB2251	—	SA2251	—
	100	SB2261	SB3261	SA2261	SA3261
	150	SB2271	SB3271	SA2271	SA3271
4.6	10	SB2411	SB3411	SA2411	SA3411
	50	SB2441	SB3441	SA2441	SA3441
	75	SB2451	—	SA2451	—
	100	SB2461	SB3461	SA2461	SA3461
	150	SB2471	SB3471	SA2471	SA3471
	250	—	SB3481	—	SA3481
10.0	250	—	SB3781	—	SA3781
20.0	250	—	SB3881	—	SA3881

